



Core Concepts: Antibacterial Drugs II Gram Positive Organisms

Helen W. Boucher, MD, FACP, FIDSA, (Hon) FRCPI
Dean and Professor of Medicine
Tufts University School of Medicine
Chief Academic Officer, Tufts Medicine

8/4/2025

1



Disclosures of Financial Relationships with Relevant Commercial Interests

Editor

- ID Clinics of North America
- Antimicrobial Agents and Chemotherapy
- Sanford Guide

2

Overview of Antibacterial Mechanisms

To Orient You: Little is Testable




Figure 20-14 Brock Biology of Microorganisms 11/e
© 2006 Pearson Prentice Hall, Inc.

3

3

Cell Wall Active Agents

- Penicillins
- Cephalosporins
- Carbapenems
- Vancomycin
- Daptomycin
- Polymyxins
- Aztreonam


4

4

12 Core Concepts: Antibacterial Drugs II Gram Positive Organisms

Helen W. Boucher, MD, FACP, FIDSA, (Hon) FRCPI

β-lactam Spectrum

- Penicillins
 - Semi-synthetic penicillins
 - 1st gen cephalosporins
 - 2nd gen cephalosporins
 - 3rd gen cephalosporins
 - 4th gen cephalosporins
 - Carbapenems
 - Monobactams
- 
- Gram-positive
- Gram-negative

5

β-lactam Antibiotics Share Mechanism of Action

— Why are there different spectrum of activity for penicillins, cephalosporins, carbapenems?

- Broad and narrow susceptibility to beta-lactamases
- Different penicillin binding proteins
- Selective efflux pumps
- Ability to reach target site

6

β-lactam Adverse Effects

- Anaphylaxis / allergy
 - See lecture by Dr. Sandy Nelson
- Seizures
 - Imipenem, cefepime
- Myelosuppression, leukopenia, hemolytic anemia
- Hypersensitivity hepatitis: e.g., Oxacillin
- Biliary stasis/sludging
 - Ceftriaxone
- Renal
 - Interstitial nephritis

7

Question #1

What is the only cephalosporin active against MRSA?

- A. Cefpodoxime
- B. Cefapime
- C. Ceftaroline
- D. Cefixime
- E. Cefoxitin

8

7

8

Cephalosporins

- Bactericidal
 - inhibit bacterial cell wall synthesis
- Time dependent killing
- Resistance mostly due to susceptibility to β -lactamases
- Fewer allergic reactions than PCN
- CSF penetration with third generation
- Most renally excreted

9

9

Key Points About Cephalosporin Activity

- Enterococci
 - None are active
- MRSA
 - Only ceftaroline and ceftobiprole active
- Anaerobic activity
 - Only Cephamycins active
 - (e.g., cefoxitin, cefotetan)
 - Now high levels of resistance

10

10

Ceftaroline Fosamil – a Prodrug (IV and IM, Not Oral)

- Activity
 - Gram-positive including MRSA and MDR *S. pneumoniae*
 - Some activity vs *E. faecalis*; not *E. faecium*
 - Limited activity vs. anaerobes
 - Active vs *Cutibacterium* (formerly *Propionibacterium*) *acnes*, *Actinomyces* spp.

Lodise & Low, Drugs, 2012; Saravolatz et al. CID 2011; 52: 1156

11

11

Ceftaroline Fosamil – a Prodrug (IV and IM, Not Oral)

- Activity
 - Active vs Gram-negative pathogens
 - *E. coli*, *Klebsiella* spp., *H. influenzae* (incl B-lactamase positive), *M. catarrhalis*
 - **Not *Pseudomonas* or ESBL+ GNB**
 - **Similar spectrum to ceftriaxone**
 - Bactericidal, time dependent killing

Lodise & Low, Drugs, 2012; Saravolatz et al. CID 2011; 52: 1156


12

12

12 Core Concepts: Antibacterial Drugs II Gram Positive Organisms

Helen W. Boucher, MD, FACP, FIDSA, (Hon) FRCPI

Ceftobiprole



- Advanced spectrum IV cephalosporin
- Broad spectrum (similar to ceftaroline)
 - Active vs G+ incl PRSP, MRSA, anaerobes
 - Some activity vs *E. faecalis*; not *E. faecium*
 - Active vs Enterobacteriaceae
 - Not active vs CRE, *P aeruginosa*

Overcash et al. CID 2021; 73: e1507; Awad SS et al. Clin Infect Dis. 2014;59(1):51–61
Holland et al. N Engl J Med 2023; 389(15):1390

13

13

Ceftobiprole

- FDA approved:
 - ABSSSI, CABP, *S. aureus* bloodstream infection/Right-sided endocarditis
- Not FDA approved for VABP
 - EU approved HAP not VAP
 - Early HABP studies failed; low ceftobiprole levels found in young ICU patients
- Dosing
 - ABSSSI and CABP:
 - 667 mg IV q 8h x 5-14 days
 - *S. aureus* bloodstream infection:
 - 667mg IV q 6h day 1-8 then 667 mg IV q 8h day 9+ (thru 42)

Overcash et al. CID 2021; 73: e1507; Awad SS et al. Clin Infect Dis. 2014;59(1):51–61
Holland et al. N Engl J Med 2023; 389(15):1390

14

14

Vancomycin

- Bactericidal (slowly)
 - Inhibits bacterial cell wall synthesis
- Active against:
 - Gram-Positive Aerobes
 - Streptococcus
 - Staphylococcus
 - Enterococcus
 - Gram-Positive Anaerobes
 - Clostridia
 - Propionibacteria
 - Peptostreptococci
 - Actinomyces

15

15

Vancomycin Resistance



- VISA
 - Thick walls, generous binding sites...
- Vancomycin resistance
 - Not in Streptococcus
 - RARE in Staphylococcus
 - Common in Enterococcus
 - Rare in *E. faecalis* (4% in 2014)
 - Common in *E. faecium* (71% in 2014)
 - Mechanism
 - Change in vancomycin binding site on peptidoglycan

16

16

Vancomycin Resistance

- VISA thickened cell wall + xs vancomycin binding sites (D-Ala-D-Ala); result: vanco trapping with reduced cellular targets
- VRE – replacement of D-Ala-D-Ala with D-alanyl-D-lactate termini – result: decreased **vancomycin** binding affinity --- high level resistance: MIC increase x 1000



17

Vancomycin for MRSA Bloodstream Infection

- Controversy re: optimal therapy – see Dr. Chambers lecture
- Vancomycin trough only monitoring no longer recommended
 - Target AUC/MIC_{BMD} ratio of 400 to 600
 - (assume vancomycin MIC_{BMD} = 1 mg/L)
- Loading dose for seriously ill adults
 - 20–35 mg/kg can be considered
 - Pediatric doses higher
 - 60-80 mg/kg/day divided q 6-8 hours

Dosing Calculator helps!

<https://www.idsociety.org/practice-guideline/vancomycin/>



18

Vancomycin ADRs / Interactions

Adverse Drug Reactions

- Nephrotoxicity
 - Duration > 14d
 - Dose > 4g / day
 - Trough > 20
- Ototoxicity
- Histamine Release Syndrome
- DRESS
- Immune thrombocytopenia
- Neutropenia
- IgA bullous dermatitis



Drug Interactions

- Increased nephrotoxicity when given with other nephrotoxins
 - Aminoglycosides
 - NSAIDs
 - Contrast
 - Cyclosporine
 - Tacrolimus
 - Loop Diuretics
 - ACE inhibitors
 - Pip/tazo (pseudo interaction)

19

19

Daptomycin (IV)

- Antimicrobial Class: Lipopeptide
- Broad spectrum gram + activity
 - Including MRSA
- Rapidly bactericidal
- Concentration-dependent killing
- Indications
 - cSSSI
 - *S. aureus* bloodstream infection
 - Right-sided endocarditis

Fenton C et al. *Drugs* 2004; 64: 445-55, Tedesco KL, Rybak MJ. *Pharmacother* 2004; 24:41-57, Mangili A et al. *Clin Infect Dis* 2005; 40:1058-60, Fowler VG et al. *New Engl J Med* 2006; 355:653-665

20

20

Daptomycin for *S. aureus* Bacteremia and Right IE

- **Pneumonia**
 - Do not use: surfactant binding inactivates drug
- **Monitoring**
 - CPK twice weekly
 - Discontinue if myopathy or CPK > 5x ULN
- **Toxicity**
 - Eosinophilic Pneumonia
 - Rx supportive care and steroids
 - Falsely prolonged Prothrombin Time
 - Muscle inflammation
 - CPK increase, myopathy, myositis
 - Risk factors: renal failure, statins, obesity

21

21

Vancomycin and Daptomycin

Drug	Mechanism of Action	Mechanism of Resistance	Spectrum	Adverse Event
Vancomycin	Inhibits cell wall synthesis (not a beta lactam)	Change in cell wall terminus from D-ala-D-ala to D-ala-D-lactate (high level resistance)	Gram positive cocci only including MRSA	<ul style="list-style-type: none"> • Histamine release syndrome • Kidney toxicity
Daptomycin	Cell membrane depolarization Potassium efflux	<ul style="list-style-type: none"> • Decreased binding of drug to cell membrane • Altered cell membrane potential 	Resistant gram positive cocci including MRSA and VRE Inactivated by surfactant (not for pneumonia)	<ul style="list-style-type: none"> • Skeletal muscle toxicity

22

22

Oritavancin and Dalbavancin Long Acting Glycopeptides

- **Mechanism of Action**
 - Similar to vancomycin
 - Inhibition of cell wall synthesis
- **Dosing**
 - Oritavancin: IV only: 1 dose (1200 mg over 3hours)
 - Dalbavancin: IV only: 1000mg, then 500mg every 7 days ...OR 1500mg x 1
- **Approved**
 - Skin and Soft Tissue
 - Oritavancin FDA warning against use in osteomyelitis
 - Dalbavancin also used for osteomyelitis, right sided endocarditis
- **Toxicity**
 - Oritavancin prolongs aPTT (artificially), PT, and activated whole blood clotting time (ACT) for 5 days

23

23

Lipo/glycopeptide Testable Toxicities

- **Vancomycin:** Nephrotoxicity, Histamine Release
- **Daptomycin:** CPK elevation, myopathy, rhabdomyolysis; eosinophilic pneumonia
- **Telavancin:** Nephrotoxicity
- **Oritavancin:** LFT elevation, false prolongation of aPTT
- **Dalbavancin:** LFT elevation

24

24

Question #2

PREVIEW QUESTION

INFECTIOUS
DISEASE
BOARD REVIEW
2025



Which quinolone has activity against MRSA?

- A. Ciprofloxacin
- B. Moxifloxacin
- C. Trovafloxacin
- D. Delafloxacin
- E. Levofloxacin

25

25

Antibiotics Active Intracellularly

- Fluoroquinolones
- Tetracyclines
- Linezolid
- TMP/SMX
- Pleuromutilins
- Linezolid/Tedizolid

26

26

Fluoroquinolone Mechanism of Action and Resistance

- Topoisomerase inhibitors
 - Inhibits DNA gyrase and topoisomerases II and IV
 - Gyrase more for gram negs, topos for gram pos
- Resistance
 - Target site mutations
 - Drug permeability mutations
 - Occurs spontaneously on therapy
 - Susceptible to drug modifying enzymes

27

27

Fluoroquinolones: Spectrum of Gram-Positive Activity

	Gram-positive	Gram-negative	Anaerobes
Cipro	Poor strep Some MSSA	Best FQ for •Pseudomonas •E coli	Some
Levo	Good strep Some MSSA	Best for <i>Stenotrophomonas</i> spp.	Some
Moxi	Good strep Good MSSA	Not effective Don't use for UTI	Best

Drs. Tamma and Gilbert will address Gram-negative activity

28

28

12 Core Concepts: Antibacterial Drugs II Gram Positive Organisms

Helen W. Boucher, MD, FACP, FIDSA, (Hon) FRCPI

Fluoroquinolone Pharmacokinetics

- High oral bioavailability
 - >95% for moxi / levo, 70-80% for cipro
 - Potential low bioavailability when taken with multivalent cations – chelation blocks absorption
- Widely distributed to tissues
 - Lower than serum but therapeutic concentration in CSF, saliva, bone, ascitic fluid and prostate gland
- Elimination
 - Levo / cipro: renal through tubular secretion
 - Moxi: >60% hepatic/ biliary unchanged

29

29

Fluoroquinolone Adverse Effects

- *C. difficile*
- Arthropathy/cartilage toxicity / tendonitis
 - FDA Warning for rare tendon rupture
 - Increased risk: advanced age, poor renal function, concomitant steroids
- Altered mental status (HA, dizziness, insomnia)
- Dysglycemia-FDA warning especially for older adults and diabetics
 - Hypo- and hyperglycemia
- Aortic aneurysm and aortic dissection-FDA warning
 - Association is controversial
- QTc Prolongation:
 - Moxi > levo ? Cipro
 - Increased risk:
 - Concomitant QTc prolongers, cardiomyopathy, bradycardia, low K⁺ and Mg⁺⁺

30

30

Delafloxacin

- Broad spectrum fluoroquinolone
- Potential advantages:
 - MRSA activity
 - Broad spectrum including *Pseudomonas*
- Dosing IV and oral twice daily
- Approved for skin and soft tissue infections

Saravolatz LD and Stein GE. Clin Infect Dis. 2019;68(6):1058–62

31

31

Tetracyclines: Major Clinical Uses

- Acne (minocycline)
- Respiratory tract infections
 - Atypical pneumonia
- Sexually Transmitted Diseases
 - Syphilis (*T. pallidum*) – alternative therapy
 - *Chlamydia* spp.
- Tick-Borne Illnesses
 - Lyme disease
 - Anaplasmosis
 - Ehrlichiosis
 - Rocky Mountain Spotted Fever
- Community Acquired MRSA infections

32

32

Tetracyclines: Adverse Effects

- Gastrointestinal
 - Nausea
 - Esophageal ulceration
 - Hepatotoxicity
- Skin
 - Photosensitivity
- Children
 - Yellow brown tooth discoloration if age <8 yrs for tetracyclines
 - **Doxycycline therapy OK for ≤21 days in children of all ages**
 - Ref: Redbook 2018 and Am Academy Pediatrics
- Pregnancy
 - Tetracyclines cross the placenta; accumulate in fetal bone/teeth
 - Most tetracyclines contraindicated in pregnancy

33

33

Newer Tetracyclines

	Omadacycline	Eravacycline
FDA approval	ABSSSI, CABP	cIAI, not cUTI (failed studies)
Dosing	200 mg loading dose over 60 min day 1, 100mg IV over 30 min or 300mg orally once daily No dose adjustment for renal/hepatic impairment	1mg/kg IV q 12h (over 60 minutes) Dose adjustment with hepatic impairment
Activity	Broad spectrum: Gram-pos including MRSA, VRE; Gram-neg including ESBL, CRE (not all); anaerobes	
Issues	Limited activity vs carbapenem-resistant <i>K. pneumoniae</i>	High MIC <i>Pseudomonas</i> , <i>Burkholderia</i> spp.
Safety	GI, rash, ?heart rate	GI, rash

34

34

Question #3

PREVIEW QUESTION

2025
INFECTION
DISEASE
BOARD REVIEW



What is the major advantage of tedizolid compared to linezolid?

- Longer half life
- Better penetration of prostate
- Better CSF Penetration
- Wide spectrum of activity against anaerobes
- More effective in clinical studies for VRE

35

35

Linezolid and Tedizolid: Oxazolidinone Drug Class

- Mechanism
 - Binds 50s ribosome/prevents formation of initiation complex
- Spectrum of activity
 - Gram positive cocci including MRSA and VRE
 - Linezolid resistant *S.aureus* reported
 - Mycobacteria
- Resistance is rare; target change
- Linezolid twice daily; Tedizolid once daily
- FDA approvals for Linezolid:
 - Skin and Soft Tissue, Pneumonia, VRE
 - NOT Bloodstream infection (Black Box Warning)

Shinabarger DL et al. Antimicrob Agents Chemother 1997; 41: 2132-36; Swaney Sm et al. Antimicrob Agents Chemother 1998; 42: 3251-55; French G. Int J Clin Pract 2001; 55: 59-63

36

36

Linezolid Adverse Events

- Adverse events related to mitochondrial toxicity:
 - Cytopenias
 - Monitor CBC
 - Peripheral and irreversible optic neuropathy
 - Rare:
 - Lactic acidosis, **serotonin syndrome (w SSRIs)**
- ↑ mortality in study of intravenous catheter-associated bacteremia

Tsioltras S et al. *Lancet* 2001;358: 207-208; Pillai SK et al. *Clin Infect Dis* 2002; 186: 1603-7; Wilson P et al. *J Antimicrob Chemother* 2003;51:186-88; Medwatch March 16, 2007

37

37

TMP/SMX Spectrum of Activity - Typical Bugs

- Gram Positive
 - Staphylococci: great
 - Streptococci: controversial
 - Enterococcus: not effective
- Gram Negative
 - *E. coli*: ok, increasing resistance
 - Enterobacterales: relatively effective
 - Pseudomonas / Acinetobacter: not effective
 - Stenotrophomonas: often drug of choice (2024 IDSA Guidance suggests combination with cefiderocol, minocycline or levofloxacin)

38

38

TMP/SMX Spectrum of Activity - Odd Bugs

- *Stenotrophomonas maltophilia*
- *Listeria monocytogenes*
- *Nocardia*
- *Moraxella catarrhalis*
- *Pneumocystis jirovecii*
- *Toxoplasmosis gondii* (but not superior to pyr/sulf)
- *Chlamydia* (but enough resistance that its not used for STDs)
- Atypical mycobacteria

39

39

Lefamulin

- Pleuromutilin antibiotic with IV and PO formulation
 - Protein synthesis inhibitor
 - Bacteriostatic
- FDA Approved community acquired bacterial pneumonia
 - Non-inferior to moxifloxacin for CABP in two studies
 - **5 days** of po lefamulin vs. 7 days of po moxifloxacin

File CID 2019

40

40

Macrolides (Erythro, Clarithro, Azithro)

Protein Synthesis Inhibitor Binds 50s Ribosome

Spectrum:

CABP Pathogens:

- *Streptococcus pneumoniae*
- *Haemophilus influenzae*
- *Moraxella catarrhalis*
- *Legionella* spp.
- *C. pneumoniae*
- Streptococcus groups A, C, and G

Strep Pneumo Resistance

- Rising rates in US
 - Don't use macrolides if local rates of resistance > 25%

41

41

Macrolide Spectrum

STDs

- *Haemophilus ducreyi* (chancroid)
- *Chlamydia* spp.

GI pathogens

- *Campylobacter* spp.
- *Helicobacter pylori*
- *Salmonella typhi*
- *Shigella* spp.

Miscellaneous Bugs

- *Arcanobacter* spp.
- *Bartonella henselae* (cat-scratch)
- *Bordetella pertussis*
- Atypical mycobacteria
- *Borrelia burgdorferi*
- *Babesia microti*

42

42

Macrolide Adverse Drug Reactions

- QTc Prolongation
 - Ery ≥ clarith > azith
- GI intolerance: nausea, bloating, diarrhea
 - Ery >> clarith >> azith
 - Dose related
 - Activity at motilin (peristalsis) receptors
 - Rare cholestatic hepatitis
- Pregnancy risk

43

43

Clindamycin Adverse Events

- Allergic reactions:
 - Rash, fever, erythema multiforme, anaphylaxis
- Elevated AST/ALT
 - Rare progression to severe liver injury
- Diarrhea
 - Can cause severe *C. difficile* toxin-mediated colitis
- Reversible neutropenia, thrombocytopenia, and eosinophilia
- Taste disturbance

Sanford Guide, Brit J Clin Pharmacol 64:542, 2007; Clin Med Insights Case Rep 2019 Dec 25;12:1-4

44

44

Thank You!


- Henry Masur
- Sue Cammarata
- G. Ralph Corey
- Sara Cosgrove
- Mike Dudley
- Mike Dunne
- **David Gilbert**
- Susan Hadley
- Teena Kohli
- Kenneth Lawrence
- Evan Loh
- Paul McGovern
- Federico Perez
- Debra Poutsika
- George H. Talbot
- Our patients and their families

45

45

Questions, Comments?

- @hboucher3
- Helen.boucher@tufts.edu
- Helen.boucher@tuftsmedicine.org



46

46