



31 Core Concepts: Antifungal Drugs
Speaker: Taison Bell, MD, MBA, FIDSA



Syndromes in the ICU that Infectious Disease Physicians Should Know

Taison Bell, MD, MBA, FIDSA
Interim Chair, UVA Department of Medicine
Division of Pulmonary and Critical Care Medicine
Division of Infectious Diseases and International Health

7/11/2025

1



Disclosures of Financial Relationships with Relevant Commercial Interests

- List of disclosures or “None”

2

Question #1

What proportion of patients in the ICU develop fever during their stay?

- A. Less than 5%
- B. Between 15-25%
- C. Over 50%
- D. Everyone. Absolutely everyone.

3

Question #1

What proportion of patients in the ICU develop fever during their stay?

- A. Less than 5%
- B. Between 15-25%
- C. Over 50%
- D. Everyone. Absolutely everyone.

4

Exam Blueprint: Critical Care Topics ~8-10%

Critical Care Medicine

- Systemic inflammatory response syndrome (SIRS) and sepsis
- Ventilator-associated pneumonias
- Noninfectious pneumonias (eosinophilic and acute respiratory distress syndrome [ARDS])
- Bacterial pneumonias Viral pneumonias
- Hyperthermia and hypothermia
- E-cigarette or vaping product use-associated lung injury (EVALI)

Medical Content Category	% of Exam
Bacterial Diseases	27%
Human Immunodeficiency Virus (HIV) Infection	15%
Antimicrobial Therapy	9%
Viral Diseases	7%
Travel and Tropical Medicine	5%
Fungi	5%
Immunocompromised Host (Non-HIV Infection)	5%
Vaccinations	4%
Infection Prevention and Control	5%
Internal Medicine and Non-Infectious Syndromes	18%
	100%

5

Question #2

- You are asked to see a 35-year-old woman with a history of seizure disorder admitted to the ICU with a fever to 40°C, hypotension, and a maculopapular rash
- She is being empirically treated with vancomycin and piperacillin-tazobactam. Blood, urine, and sputum cultures (taken prior to antibiotic initiation) are negative
- Exam: Tachycardia with otherwise normal vital signs. Diffuse maculopapular rash with facial edema and sparing of the mucosal surfaces
- Labs are notable for elevated AST/ALT and peripheral eosinophilia
- Only home medication is lamotrigine, which was started two weeks prior to admission

6

Question #2

What is her clinical syndrome is most consistent with?

- A. Sepsis
- B. Stevens–Johnson syndrome (SJS)/toxic epidermal necrolysis (TEN)
- C. DRESS (drug-induced hypersensitivity syndrome)
- D. Erythema Multiforme
- E. Neuroleptic Malignant Syndrome (NMS)

7

Question #2

What is her clinical syndrome is most consistent with?

- A. Sepsis
- B. Stevens–Johnson syndrome (SJS)/toxic epidermal necrolysis (TEN)
- C. DRESS (drug-induced hypersensitivity syndrome)
- D. Erythema Multiforme
- E. Neuroleptic Malignant Syndrome (NMS)

8

Morbilliform Rash with Facial Edema and Eosinophilia



9

DRESS (Drug-induced Hypersensitivity Syndrome)

Rash Characteristics	Morbilliform involving >50% BSA, inflamed, facial edema, infrequent mucosal involvement
Onset	Usually 1-3 (up to 6) weeks after drug exposure
Other Features	Fever, LAD, other organ involvement in 80% (liver, kidney, pancreas, heart, lung), expansion of CD4/8 T cells → Herpesviridae reactivation (HHV6)
Lab Findings	Eosinophilia, lymphocytosis/lymphopenia, atypical lymphocytes
Classic Meds	Aromatic AEDs (highest with lamotrigine), Vancomycin, Raltegravir, Dapsone and other Sulfas, anti-TB RIPE
DDx	SLE, mycoplasma, viral hepatitis, mononucleosis
Treatment	Withhold offending agent, supportive care Steroids, CsA, IVIg are controversial. Mortality is high

10

Exanthematous Drug Eruptions

- T-cell-mediated, delayed type IV hypersensitivity reaction
- Diffuse maculopapular rash (morbilliform)
- Highest incidence with aromatic antiseizure medications: carbamazepine, phenytoin, and lamotrigine (1:100)

SJS/TEN	AGEP	DRESS
<ul style="list-style-type: none">• Severe blistering• Mucosal involvement common• SJS: <10% BSA• TEN: >30% BSA	<ul style="list-style-type: none">• Rapidly spreading (hours) pustular lesions• Mucosal involvement rare• Common ddx: psoriasis	<ul style="list-style-type: none">• > 50% BSA• Facial edema• Infrequent mucosal involvement• Eosinophilia

11

Stevens Johnson Syndrome and Toxic Epidermal Necrolysis

Rash Characteristics	Erosive mucositis of oral, urogenital, and ocular sites
Onset	4-28 days after drug exposure
Other Features	Fever, partial or full thickness injury with painful necrolysis, pulmonary and GI manifestations
Lab Findings	Leukopenia, no eosinophilia
Risk Factors	Aromatic AEDs, infection (mycoplasma), GVHD, HIV
Treatment	Withhold offending agent, supportive care Steroids and IVIg are controversial

12

Stevens Johnson Syndrome and Toxic Epidermal Necrolysis




- “Positive Nikolsky sign”
 - Slight rubbing of the skin results in exfoliation of the outermost layer
 - NOT specific for Stevens Johnson and TEN
 - Staph scalded skin syndrome (mostly children, no mucosal involvement)
 - Pemphigus
 - Others

13

Erythema Multiforme

- Immune mediated
- Distinctive target lesions that are usually asymptomatic
 - Febrile prodrome in some cases
- Often associated with oral, ocular, and genital mucosal lesions
- Less severe than DRESS or SJS or TEN
- Causes: Infection > Drugs
 - Infections: HSV, Mycoplasma, many others
 - Cancer, autoimmune, drugs, etc.
- Self Limiting in 10-14 days



14

Extreme Hyperpyrexia (T>41.5C)

Heat Stroke

- Exertional (football player in August)
- Non-exertional (Elderly)
- Lack of hydration and/or inability to sweat

Drugs

- Cocaine, ecstasy etc.

The Pyrexia Syndromes

15

Question #3

- You are called to the PACU to see a 29-year-old previously healthy male with a fever of 41.6°C who is 4 hours post-op from an arthroscopy for a rotator cuff injury
- He initially did well post operatively except for some nausea that was treated
- The patient is somnolent, flushed, diaphoretic, and rigid. His blood pressure has risen from 130/70 to 180/100 but is now dropping. He is given one ampule of Narcan but does not respond

Which of the following would you give?

- A. Antihistamines
- B. High-dose corticosteroids
- C. Dantrolene
- D. IVIG
- E. Dilantin

16

Question #3

- You are called to the PACU to see a 29-year-old previously healthy male with a fever of 41.6°C who is 4 hours post-op from an arthroscopy for a rotator cuff injury
- He initially did well post operatively except for some nausea that was treated
- The patient is somnolent, flushed, diaphoretic, and rigid. His blood pressure has risen from 130/70 to 180/100 but is now dropping. He is given one ampule of Narcan but does not respond

Which of the following would you give?

- A. Antihistamines
- B. High-dose corticosteroids
- C. Dantrolene
- D. IVIG
- E. Dilantin

17

Malignant Hyperthermia

- Syndrome - Rare (~700 cases/year) but 5-10% mortality
 - Early signs: Steep rise in CO2, tachycardia, tachypnea, muscle rigidity/contraction (masseter spasm)
 - Late signs: Hyperthermia, acidosis, hyperkalemia, cardiac arrhythmias
- Genetic defect in the RYR1 or (less commonly) CACNAS1S gene
 - Ca++ transport in skeletal muscle
 - Autosomal dominant
 - (excessive calcium accumulation)
- Triggers
 - Usually < 1 hour after trigger (up to 10 hours)
 - Classic: Volatile anesthetics (halothane, sevoflurane, desflurane), succinylcholine

18

Neuroleptic Malignant Syndrome (NMS)

- Frequent trigger = haloperidol
 - Any "neuroleptic" (antipsychotic)
 - Lead pipe rigidity
 - Antiemetics such as metoclopramide
 - Withdrawal of antiparkinson drugs (L dopa)
- Onset variable: 1-3 days/within first 2 weeks
 - Time of drug initiation
 - When dose changed
- Management
 - Dantrolene
 - (Direct muscle relaxant for up to 10 days)
 - Dopamine agonists (bromocriptine and others)

19

Serotonin Syndrome

Clinical Characteristics of Serotonin Syndrome	
Pathogenesis	Excess Serotonergic Activity <ul style="list-style-type: none">• Therapeutic drugs, drug interactions, self poisoning
Triggers	<ul style="list-style-type: none">• Linezolid = MAO Inhibitor• SSRI inhibitors (Bupropion)• Antiemetics (Granisetron)• Tricyclic antidepressants (amitriptyline)
Clinical Manifestations	<ul style="list-style-type: none">• Acute onset (within 24 hrs of new drug/drug change)• Hyper-reflexive>bradyreflexia• Nausea, vomiting, diarrhea, tremors followed by shivering
Treatment	<ul style="list-style-type: none">• Withdraw offending medication• Consider benzodiazepines and cyproheptadine

20

What to Look for on the Exam

	Malignant Hyperthermia	NMS	Serotonin Syndrome
Trigger	Succinylcholine or inhaled halogenated anesthesia	Withdrawal of L Dopa in Parkinsons or Neuroleptic Drugs	SSRIs, Antiemetics, Linezolid, Lithium, Street Drugs
Onset	Rapid onset in perioperative period	Subacute over 1-3 days	6-24 hours of starting a drug or increasing dose
Exam	Masseter spasm, Lead pipe rigidity	Mental status change with dysautonomia, catatonia, mutism, stupor, coma	Shivering, myoclonus, n/v/d, hyper-reflexia, flush skin
Labs	Severe hypercarbia, rhabdomyolysis	CK rise, myoglobinemia	Nothing classic

21

Hypothermia: <35°C

- Causative Drugs
 - Beta blockers (metoprolol)
 - Alpha blockers (clonidine)
 - Opioids
 - Ethanol
 - Syndrome
 - Hypotension due to fluid shifts
 - ***Give broad spectrum antibiotics empirically if they fail to raise temperature 0.67°C/hour**
 - Consider adrenal or thyroid insufficiency
 - Treatment
 - Rewarming
 - "ABC"s
 - o Airway, Breathing, Circulation
- Antidepressants
Antipsychotics
Aspirin
Oral hypoglycemics

22

Question #4

- You are called to the medical ICU to see a 47 y/o woman with a history of alcoholic cirrhosis with ARDS and shock
- Initially admitted to general medicine for encephalopathy in the setting of skipping lactulose doses
- On HD#3 developed ARDS, thought to be from aspiration
- Subsequently goes into distributive shock. Started on vancomycin and piperacillin-tazobactam
- Patient has daily fevers to 39°C and a persistent low-dose levophed requirement
- Labs: mild hyponatremia and hyperkalemia. Metabolic acidosis
- Micro: blood, urine, sputum, and ascitic fluid are benign
- Radiology: CXR with unchanged b/l multifocal opacities, RUQ USG benign, Abd CT benign

23

Question #4

Which of the following would you give?

- A. Broader spectrum antibacterial treatment
- B. Stress dose corticosteroids
- C. Dantrolene
- D. IVIG
- E. Antifungal therapy

24

Question #4


Which of the following would you give?

- A. Broader spectrum antibacterial treatment
- B. Stress dose corticosteroids**
- C. Dantrolene
- D. IVIG
- E. Antifungal therapy

25

Differential Diagnosis of Shock

Ohm's Law \longrightarrow




26

Why not Empiric Antifungal? EMPIRICUS

Multi-center RCT of 260 Adults in ICU

- Non-neutropenic
- Multiorgan failure
- ICU-acquired sepsis
- On MV at least 5d
- At least 4d broad spectrum Abx in prior week
- Multifocal candida colonization



27

Question #5



A patient with end stage renal disease on dialysis through a tunneled hemodialysis catheter is admitted to the medical ICU with altered mental status, hypotension, and fever. On exam he has obvious purulence at the catheter site.

For the patient's syndrome, which of the following is NOT an evidence-based intervention?

- A. Early and effective antibiotics
- B. Albumin as the preferred resuscitation fluid
- C. Measuring serum lactate
- D. Fluid resuscitation with 30 cc's/kg crystalloid

28

Question #5

PREVIEW QUESTION

OVER
INFECTIOUS
DISEASE
BOARD REVIEW
2025



A patient with end stage renal disease on dialysis through a tunneled hemodialysis catheter is admitted to the medical ICU with altered mental status, hypotension, and fever. On exam he has obvious purulence at the catheter site.

For the patient's syndrome, which of the following is NOT an evidence-based intervention?

- A. Early and effective antibiotics
- B. Albumin as the preferred resuscitation fluid**
- C. Measuring serum lactate
- D. Fluid resuscitation with 30 cc's/kg crystalloid

29

FYI: Sepsis 3 Definition: Not Testable!

- Definition of Sepsis
 - “Life-threatening organ dysfunction due to a dysregulated host response to infection”
- Definition of Septic Shock: Sepsis
 - Absence of hypovolemia
 - Vasopressor to maintain mean blood pressure >65mmg
 - Lactate >2 mmol/L (>18 mg/dL)
- Predicting Outcome
 - Increase in the Sequential Organ Failure Assessment (SOFA) score (10% mortality)
 - Quick Sofa is relatively specific but not very sensitive

30

Sepsis 3 Definition: For Background (Not Testable)!

	Traditional Definition	Sepsis 3
Sepsis	Suspected or known infection with ≥ 2 SIRS criteria	Life-threatening organ dysfunction due to a dysregulated host response to infection <ul style="list-style-type: none">- SOFA score ≥2 points or positive qSOFA
Severe Sepsis	Sepsis + organ failure	N/A
Septic Shock	Severe sepsis + hypotension refractory to adequate fluid resuscitation or addition of vasopressors	Sepsis with adequate resuscitation with vasopressor requirement and lactate ≥ 2 mmol/L

Increase in the Sequential Organ Failure Assessment (SOFA) score (10% mortality)
Quick Sofa is relatively specific but not very sensitive

31

Surviving Sepsis Campaign

Managing Sepsis

What's the Bottom Line?


- Some recommendations are plausible
 - Fluid resuscitation with 30 cc's/kg crystalloid
 - Vasopressors for MAP goal 65
 - But do not use Dopamine!
- Some are wrong
 - Early goal directed therapy
 - Tight glucose control. Better outcomes <180
- Two are unequivocally true
 - Early effective antibiotics
 - Source control

32

31 Core Concepts: Antifungal Drugs
Speaker: Taison Bell, MD, MBA, FIDSA

Surviving Sepsis Campaign

Other Things



Stress-dose steroids: conflicting data


- CORTICUS/ADRENAL
 - No change in mortality with hydrocortisone
 - **Quicker reversal of shock**
- Annane/APROCCHSS
 - Improved mortality with hydrocort/fludricort
 - **Quicker reversal of shock**

Antiendotoxin and Anticytokine therapy

- No benefit

Antithrombosis (Activated Protein C)

- Taken off the market





33

Surviving Sepsis Campaign Bundles

3 Hour Bundle	6 Hour Bundle
<ul style="list-style-type: none">- Measure lactate level- Draw blood cultures- Administer broad spectrum antibiotics- Administer 30 cc/kg IV crystalloid	<ul style="list-style-type: none">- Start vasopressors if MAP <65 despite fluid resuscitation- Reassess volume status if hypotension persists after fluid resuscitation or if initial lactate ≥ mmol/L

34

Ventilator Associated Pneumonia



35

Ventilator Associated Pneumonia
National Healthcare Safety Network

Pathogen	% of Isolates
Staph aureus	24.7%
Pseudomonas aeruginosa	16.5%
Klebsiella	10%
Enterobacter	8. %
E. Coli	5%

36

IDSA VAP Treatment Guidelines

Cover for *S. aureus*, *P. aeruginosa*, and other GNRs in ALL patients (strong recommendation, very low-quality evidence)

Clinical Question	Recommendation
MRSA coverage	Use vancomycin or linezolid
PsA and other GNRs	Pip-tazo, Cefepime, Ceftazidime, Levofloxacin
Double GNR coverage?	Only if >10% of isolates are resistant to the primary abx
Double coverage agent	FQs, aminoglycosides (no monotherapy), polymyxins
Procalcitonin	Do not use for diagnosis. Consider to aid in discontinuation
Duration of therapy	7 days, consider longer or shorter based on clinical signs

Clin Infect Dis 2016; 63: e61-e111

37

Question #6

34-year-old woman with opiate use disorder is admitted to the medical ICU for acute respiratory distress syndrome requiring intubation. She has been receiving intravenous daptomycin through a PICC for tricuspid valve endocarditis for the past three weeks. Transthoracic echo is unchanged from prior and chest CT shows bilateral ground glass opacities with scattered areas of consolidation. Blood cultures are negative. Bronchial alveolar lavage shows a predominance of eosinophils with negative cultures.

Which of the following is the most likely cause of her respiratory illness?

- A. Injection drug use
- B. Septic pulmonary emboli
- C. Daptomycin
- D. Sepsis

38

Question #6

34-year-old woman with opiate use disorder is admitted to the medical ICU for acute respiratory distress syndrome requiring intubation. She has been receiving intravenous daptomycin through a PICC for tricuspid valve endocarditis for the past three weeks. Transthoracic echo is unchanged from prior and chest CT shows bilateral ground glass opacities with scattered areas of consolidation. Blood cultures are negative. Bronchial alveolar lavage shows a predominance of eosinophils with negative cultures.

Which of the following is the most likely cause of her respiratory illness?

- A. Injection drug use
- B. Septic pulmonary emboli
- C. Daptomycin
- D. Sepsis


39

Eosinophilic Pneumonia

- Rare disorder characterized by eosinophil infiltration of the pulmonary parenchyma
- Often associated with peripheral eosinophilia
- Many drugs linked: daptomycin, nitrofurantoin, amiodarone, ACE-i's, etc.
- Daptomycin-induced EP: precise mechanism unknown but believed to be related to daptomycin binding to pulmonary surfactant leading to epithelial injury

40

ARDS Management



41

Question #7

A 22-year-old male presents to the ED with a three-week history of cough, shortness of breath, and low-grade fever. His past medical history is unremarkable. There are no sick contacts or recent travel. He went to an urgent care center one week ago and was prescribed levofloxacin but has not improved. ROS is notable for frequent use of e-cigarettes with THC-containing products. Physical examination reveals mild tachycardia, tachypnea, and decreased breath sounds bilaterally. His oxygen saturation is 88% on room air. A chest X-ray shows bilateral diffuse opacities. Laboratory studies reveal an elevated white blood cell count and elevated inflammatory markers.

42

Question #7

What is the most likely diagnosis?

- A. E-cigarette or vaping product use–associated lung injury (EVALI)
- B. Community acquired pneumonia
- C. Acute respiratory distress syndrome (ARDS)
- D. Tuberculosis
- E. Pulmonary embolism

43

Question #7


What is the most likely diagnosis?

- A. E-cigarette or vaping product use–associated lung injury (EVALI)
- B. Community acquired pneumonia
- C. Acute respiratory distress syndrome (ARDS)
- D. Tuberculosis
- E. Pulmonary embolism

44

E-cigarette or Vaping Product Use
Associated Lung Injury (EVALI)


- Primarily associated with e-cigarettes and vaping products, particularly THC-containing compounds
- Clinical presentation is very similar to community acquired pneumonia
- Exact cause not fully understood but believed to be related to direct lung injury → inflammatory response
- Treatment: supportive care, cessation of e-cigarette use



MMWR. 2019;68(46):1076

45

CLABSI



46

Antiseptic Techniques: Catheter Insertion

Hand Hygiene

- Soap & water or alcohol-based rub before/after insertion (IB)
- Sterile gloves while inserting (IA)

Skin Prep

- Chlorhexidine solution before insertion and during dressing changes (IA)
- Allow to fully dry before insertion (IB)

Barrier

- Maximum barrier protection: cap, mask, sterile gown, sterile gloves and full sterile drape (IB)

CID 2011;52 (1 May)

47

Remove the Catheter

- On the Board Exam
 - It's almost never wrong to remove/replace catheter
- Syndromes Requiring Removal
 - Septic shock
 - Septic thrombophlebitis/Venous obstruction
 - Endocarditis
 - Positive blood cultures>72 hrs after appropriate abx
- Organisms Requiring Removal
 - Staph aureus
 - Atypical mycobacteria
 - Candida species
 - Cutibacterium acnes
 - Pseudomonas aerug
 - Bacillus species
 - Malssezia
 - Micrococcus

48

Antibiotic Impregnated Catheters and Hubs Plus Antibiotic Lock Solutions

- Not likely testable on the boards
- They have a role, but not well defined

49


Near Drowning/Submersion Injuries

- Prophylactic Antibiotics
 - Not indicated unless water grossly contaminated
 - Steroids not indicated
- Etiologic Agents
 - Water borne organisms common
 - Pseudomonas, Proteus, Aeromonas
- Therapy for Pneumonia
 - Directed at identified pathogens

50

Approach

- Run med list
- Consider AI
- Pyrexia syndromes



51

Thank You

- Good luck!
- Please give feedback
- Contact
 - taison.bell@virginia.edu
 - Twitter/X: @TaisonBell (but don't bother)

52