



HIV-Associated Opportunistic Infections II

Rajesh T. Gandhi, MD
 Massachusetts General Hospital
 Professor of Medicine, Harvard Medical School
 Boston, Massachusetts

Acknowledgement: Dr. Henry Masur for slides

8/5/2025

1



Disclosures of Financial Relationships with Relevant Commercial Interests

- None

2

HIV Associated Opportunistic Infections: Part 2

- Opportunistic CNS Infections: Brain Lesions
- Opportunistic CNS Infections: Cryptococcal Meningitis
- Mycobacterial Infections
- Immune Reconstitution Inflammatory Syndrome


3

Question #1

- 50 yo M with HIV (CD4 40, HIV RNA 600,000 not on antiretroviral therapy) presents with fever, headache.
- Northeast US, no travel; no animal or TB exposures
- MRI: ring enhancing lesions; no midline shift
- Serum toxoplasma IgG +. CSF: no WBC, normal protein, toxoplasma (toxoplasma) PCR pending

What would you recommend?

- A. Brain biopsy
- B. Meningeal biopsy
- C. Initiate anti-toxo therapy; if no response in 2 weeks, brain biopsy
- D. Vancomycin, cefepime, metronidazole



4

Brain Lesions in People with HIV (PWH)




- MRI with contrast favored over CT (CT without contrast may miss lesions)
- Clues:
 - Toxoplasma: multiple ring enhancing lesions, often involving basal ganglia; serum toxoplasma IgG positive (reactivation)
 - Primary CNS lymphoma: large solitary focal brain lesion; may cross corpus callosum; increased FDG PET uptake; B cell lymphoma; CSF EBV PCR+. CD4 cell count <50
 - Tuberculoma: consider in person from endemic area with contrast enhancing lesions, basilar meningitis
 - Progressive multifocal leukoencephalopathy (PML): asymmetric non-enhancing lesions in subcortical white matter without mass effect

Siripurapu R and Ota Y, Neuroimag Clin N Am, 2023


5

Toxoplasma Encephalitis



www.idimages.org


Primary CNS Lymphoma




Siripurapu R and Ota Y, Neuroimag Clin N Am, 2023

6

PML: Asymmetric white matter changes adjacent to cortical ribbon no mass effect



HIV Encephalitis: bilateral symmetric white matter changes



www.idimages.org. Contributed by Dr. Vince Marconi

7

Evaluation of CNS Mass Lesions in People with HIV/AIDS

Toxoplasmosis
Lymphoma
Tuberculosis
Fungus
Nocardia
Bacterial
Syphilis
Kaposi
Chagoma
Glioblastoma

Radiologic Results

Non-specific although certain features suggestive
Look for Extra CNS lesions for biopsy

Laboratory Studies to Perform


Serology: Toxo IgG
Serum Cryptococcal Ag and Histoplasma Ag
Blood culture - AFB
CSF - Cryptococcal Ag
PCR (EBV, CMV, Toxoplasma, JC virus)

Response to Empiric Therapy

8

Toxoplasma Encephalitis (TE)

- Caused by protozoan, *Toxoplasma gondii*
- Reactivation of latent tissue cysts
- Highest risk is in PWH with CD4 count <100
- May present with headache, confusion, weakness, fever
- Diagnosis:
 - Serum toxoplasma IgG usually positive; negative serology makes TE unlikely
 - MRI: ring-enhancing lesions, often involving basal ganglia
 - CSF toxoplasma PCR: high specificity (96-100%); sensitivity 50-60% (negative PCR does not rule out TE)
 - Empiric diagnosis: clinical, radiographic improvement with anti-toxoplasma therapy; if no response by about 2 weeks, consider brain biopsy




<https://clinicalinfo.hiv.gov/en/guidelines/hiv-clinical-guidelines-adult-and-adolescent-opportunistic-infections/toxoplasma-gondii?view=full>

9

Time to Neurologic Response for Toxoplasma Encephalitis

35 PWH with TE Treated with Clindamycin - Pyrimethamine



10

Therapy for Toxoplasma Encephalitis


- **Preferred Regimen**
 - Sulfadiazine plus pyrimethamine plus leucovorin (PO only)
 - May be unavailable or excessively expensive
 - Trimethoprim-sulfamethoxazole (PO or IV)
 - In patients with sulfa allergy, sulfa desensitization should be attempted
- **Alternative Regimens – for those who cannot tolerate sulfonamides**
 - Clindamycin plus pyrimethamine (and leucovorin)
 - Atovaquone +/- Pyrimethamine (and leucovorin)

Note: Initiate antiretroviral therapy when patient is tolerating anti-toxoplasma therapy (usually within a week or two after starting anti-toxoplasma therapy)

<https://clinicalinfo.hiv.gov/en/guidelines/hiv-clinical-guidelines-adult-and-adolescent-opportunistic-infections/toxoplasma-gondii?view=full>

11

Compared with Sulfa-Pyrimethamine, Trim-sulfa has similar response rate, lower toxicity



12

Adjunctive Therapies for Toxoplasma Encephalitis

- Corticosteroids
 - Not routine
 - Only if mass effect, increased intracranial pressure/symptoms/signs
- Anticonvulsants
 - Should not be given prophylactically
 - Only if patients have seizures

13

Primary Prevention of Toxoplasmosis in People with HIV (PWH)

- **Indication**
 - Positive Toxoplasma IgG and CD4 <100 cells/uL
- **Drugs**
 - First Choice: TMP-SMX (one double strength tablet daily)
 - Alternatives
 - Other dosing regimens for TMP/SMX
 - Dapsone-Pyrimethamine (with leucovorin)
 - Atovaquone +/- Pyrimethamine (with leucovorin)

<https://clinicalinfo.hiv.gov/en/guidelines/hiv-clinical-guidelines-adult-and-adolescent-opportunistic-infections/toxoplasma-gondii?view=full>

14

Primary Prevention of Toxoplasmosis in PWH


- For patients with CD4<200 who are on TMP-SMX or atovaquone for PCP prophylaxis
 - Nothing more is needed
- For patient on Aerosol Pentamidine or Dapsone for PCP prophylaxis
 - If on dapsone: add pyrimethamine (plus leucovorin)
 - If on Aerosol pentamidine because cannot take TMP-SMX: not protected-
 - Consider switching to atovaquone if seropositive for toxo

<https://clinicalinfo.hiv.gov/en/guidelines/hiv-clinical-guidelines-adult-and-adolescent-opportunistic-infections/toxoplasma-gondii?view=full>


15

Case

- A 39-year-old female from Brazil presents to ED with a seizure.
 - HIV Ag/Ab is positive
 - CD4 = 20/ μ L
 - VL = 100,000 copies/ μ L
- She is started on sulfadiazine and pyrimethamine.
- After 10 days, she has not improved, and a brain biopsy is performed





16



17

Trypanosoma cruzi in Blood Smear and CSF (Chagasic Encephalitis in PWH)



Badero et al, AIDS THERAPY, 4th Ed
DiazGranados C, Lancet ID, 2009

18

HIV Associated Opportunistic Infections: Part 2

Opportunistic CNS Infections: Cryptococcal Meningitis

19

Question #2

PREVIEW QUESTION



- 50-year-old woman with HIV (CD4 20, HIV RNA 500,000) presents with fever and headache. Not on antiretroviral therapy (ART). Diagnosed with cryptococcal meningitis
- Started on induction therapy (liposomal amphotericin plus 5FC)

When should she be started on ART?

- Start ART at the same time as anti-fungal therapy
- About 4 weeks after starting anti-fungal therapy
- 6 months after starting anti-fungal therapy
- After completing a full course of maintenance anti-fungal therapy

20

HIV-Associated Cryptococcal Meningitis

- Usually presents with subacute onset of confusion, lethargy
- Neck stiffness and photophobia only occur in 25%
- May be accompanied by non-CNS manifestations: pneumonia, skin lesions, prostate infection
- CD4 Count <100 cells/uL in 90% of patients
- CSF: minimal abnormalities or lymphocytic pleocytosis with elevated protein.
- Opening pressure > 25 cm H₂O in 60-80% of patients (be sure to measure)
- Serum and CSF cryptococcal antigen positive in almost all patients.
- Blood cultures positive for cryptococcus in 60%

<https://clinicalinfo.hiv.gov/en/guidelines/hiv-clinical-guidelines-adult-and-adolescent-opportunistic-infections/cryptococcosis?view=full>

21

CLINICAL INFO HIV.gov

Enter Your Search Term...

Home About Guidelines Drug Database Glossary News Resources Contact Us Language (EN)

HOME GUIDELINES HIV CLINICAL GUIDELINES: ADULT AND ADOLESCENT OPPORTUNISTIC INFECTIONS What's New in the Guidelines

Guidelines for the Prevention and Treatment of Opportunistic Infections in Adults and Adolescents With HIV

The information in the brief version is excerpted directly from the full-text guidelines. The brief version is a compilation of the tables and boxed recommendations.

Search Guidelines Guideline Search Term...

<https://clinicalinfo.hiv.gov/en/guidelines/hiv-clinical-guidelines-adult-and-adolescent-opportunistic-infections/whats-new>

22

Therapy of Cryptococcal Meningitis

Liposomal Ampho B 3-4 mg/kg daily
plus
Flucytosine* 25 mg/kg QID

→ 2 weeks Induction

Fluconazole 800 mg po qd**

→ ≥8 weeks Consolidation

Fluconazole 200 mg po daily***

→ ≥ 52 weeks Maintenance

*5FC Associated with earlier sterilization CSF, fewer relapses, improved survival


**For clinically stable patients with negative CSF cultures, dose can be reduced to 400 mg daily

*** Stop after at least 1 yr total therapy if patient asymptomatic, CD4 >100, suppressed HIV RNA on ART


23

Single-dose Liposomal AmB with Fluconazole/5FC Preferred in resource-limited health care systems (WHO)

AMBITION Trial (n=814 participants)



All cause mortality, week 10:
No difference between groups



Adverse events less frequent in single-dose AmB group

Jarvis JN et al, NEJM, 2022

24

Management of Cryptococcal Meningitis - 1

- Patients should be followed in hospital for at least 7 days and ideally 14 days
- Lumbar puncture at day 7 and 14
- In patients with symptoms of elevated intracranial pressure and opening pressure >25 cm: remove CSF to reduce pressure by half or <20cm H2O
 - Lumbar drain or VP shunt may be needed if pressures remain elevated
- Successful induction therapy = clinical improvement and negative CSF culture
- India ink and CSF CrAg frequently positive at Week 2: not indicative of failure
- Monitoring of cryptococcal antigen titers not recommended

<https://clinicalinfo.hiv.gov/en/guidelines/hiv-clinical-guidelines-adult-and-adolescent-opportunistic-infections/cryptococcosis?view=full>

25

Management of Cryptococcal Meningitis - 2

- For flucytosine, therapeutic drug monitoring indicated.
Toxicities: marrow suppression, hepatitis, diarrhea. Renal elimination: monitor kidney function
- Not routinely recommended: Corticosteroids, Mannitol, Acetazolamide

<https://clinicalinfo.hiv.gov/en/guidelines/hiv-clinical-guidelines-adult-and-adolescent-opportunistic-infections/cryptococcosis?view=full>

26

Dexamethasone Did Not Reduce Mortality and Was Associated with More Adverse Events and Disability

ORIGINAL ARTICLE

Adjunctive Dexamethasone in HIV-Associated Cryptococcal Meningitis

J. Beardsley, M. Wolbers, F.M. Kibengo, A.-B.M. Ggayi, A. Kamali, N.T.K. Cuc, T.Q. Binh, N.V.V. Chau, J. Farrar, L. Merson, L. Phuong, G. Thwaites, N. Van Kinh, P.T. Thuy, W. Chierakul, S. Siriboon, E. Thiansukhon, S. Onsanit, W. Supphamongkhonchaikul, A.K. Chan, R. Heyderman, E. Mwinjiwa, J.J. van Oosterhout, D. Imran, H. Basri, M. Mayxay, D. Dance, P. Phimmason, S. Rattanavong, D.G. Lalloo, and J.N. Day, for the CryptoDex Investigators*


NEJM, 2016

27

When to Start ART for Cryptococcal Meningitis

- DHHS OI Guidelines recommend ART initiation 4-6 weeks after initiation of antifungal therapy
- Some experts start ART at 2-4 weeks after initiation of anti-fungal therapy with ART initiation at 2 weeks for those who have clinically improved, have control of intracranial pressure, have negative CSF cultures and can be closely monitored

COAT trial: early ART (1-2 wks) associated with higher mortality than delayed ART (5 wk)



<https://clinicalinfo.hiv.gov/en/guidelines/hiv-clinical-guidelines-adult-and-adolescent-opportunistic-infections/cryptococcosis?view=full>
Boulware D et al. NEJM. 2014
Gandhi RT et al, IAS USA Guidelines, JAMA 2024

28

Preventing Disease (Pre-emptive Therapy for Cryptococcal Ag+/Low CD4)

- Screen patients with CD4 count <200 with serum cryptococcal antigen
 - Frequency of + Ag: 2.9% if CD4 <100, 4.3% if CD4 < 50
 - Positive serum CrAg predicts development of disease
- If Positive: Perform LP and Blood Cultures to determine Rx
 - If CSF positive or serum CrAg by LFA is $\geq 1:640$: Treat like cryptococcal meningitis/disseminated (Ampho/5FC)
 - If CSF negative: fluconazole 800 to 1200 mg daily for 2 wks, then 400 to 800 mg daily for 10 wks, then 200 mg daily (total 6 months)

<https://clinicalinfo.hiv.gov/en/guidelines/hiv-clinical-guidelines-adult-and-adolescent-opportunistic-infections/cryptococcosis?view=full>

29

HIV Associated Opportunistic Infections: Part 2

Mycobacterial Infections

30


Tuberculosis in PWH: Highlights

- High risk of TB reactivation in PWH: $\approx 5\text{-}10\%$ per year; may occur even when CD4 count >200
- Screen PWH for latent TB (tuberculin skin test, TST, or IGRA); if CD4 count low, repeat TB screening after immune reconstitution on ART
- TB prophylaxis: positive TST (>5 mm) or IGRA; close contact of person with infectious TB
- When to start ART in people with HIV and TB
- CD4 count <50: start within 2 weeks of TB therapy
- CD4 count >50: start within 2-8 weeks of TB therapy (most would start sooner)
- TB Meningitis: high mortality; start ART once TB meningitis under control and at least 2 weeks after initiating TB treatment; close monitoring needed
- Prednisone may prevent paradoxical TB immune reconstitution inflammatory syndrome

<https://clinicalinfo.hiv.gov/en/guidelines/hiv-clinical-guidelines-adult-and-adolescent-opportunistic-infections/mycobacterium?view=full> Torok et al, CID, 2011; Meintjes NEJM, 2018

31

Extrapulmonary TB and High Organism Load More Common in PWH with Low CD4 Count



Jones et al, Am Rev Respir Dis, 1993; Perlman et al, CID, 1997

32

40 HIV-Associated Opportunistic Infections II

Speaker: Rajesh Gandhi, MD

Question #3

PREVIEW QUESTION



- 45-yo man with HIV (CD4 11, HIV RNA 300,000) presents with fever, diarrhea and weight loss.
- Started on dolutegravir + tenofovir/emtricitabine
- Two weeks later, develops enlarged supraclavicular lymph node
- Biopsy: necrotizing granulomas and AFB; cultures grow MAC

What would you recommend?

- A. Stop ART and initiate treatment for MAC
- B. Continue ART; initiate treatment for MAC
- C. Start steroids and stop all other treatments



Image from Riddell J, J Translational Med, 2007

33

Mycobacterium Avium Complex

- **Epidemiology**
 - Ubiquitous in the environment
- **Transmission**
 - Inhalation, ingestion
- **Risk factors**
 - CD4 <50, HIV RNA >1000
- **Clinical Manifestations of Disseminated MAC**
 - Fever, sweats, wasting, diarrhea, lymphadenopathy, hepatosplenomegaly
 - Rare as cause of lung disease
 - Labs: elevated alkaline phosphatase, anemia

<https://clinicalinfo.hiv.gov/en/guidelines/hiv-clinical-guidelines-adult-and-adolescent-opportunistic-infections/disseminated?view=full>

34

Diagnosis

- Compatible symptoms and signs along with isolation of MAC from cultures of blood, lymph node or other normally sterile sites
- MAC may be detected in respiratory or GI tract but routine screening of these sites and pre-emptive therapy for MAC is **not recommended**

<https://clinicalinfo.hiv.gov/en/guidelines/hiv-clinical-guidelines-adult-and-adolescent-opportunistic-infections/disseminated?view=full>

35

Treatment for MAC

- **Specific Therapy**
 - Clarithromycin or Azithromycin + Ethambutol
 - Rifabutin, fluoroquinolone or amikacin as a 3rd or 4th drug, particularly if severe disease ("high burden of organisms")
 - Beware drug interactions with clarithromycin or rifabutin (azithromycin has fewer drug interactions)
 - Perform susceptibility testing on MAC isolate
- **Antiretroviral Therapy**
 - Start as soon as possible after diagnosis, preferably at the same time or within a few days of initiation of anti-mycobacterial therapy

<https://clinicalinfo.hiv.gov/en/guidelines/hiv-clinical-guidelines-adult-and-adolescent-opportunistic-infections/disseminated?view=full>

36

Primary MAC Prophylaxis

- Primary prophylaxis against disseminated MAC disease is **NOT** recommended if ART initiated immediately
- People with HIV who have CD4 cell count <50, are not on ART, who remain viremic on ART or have no options for suppressive ART should receive prophylaxis after excluding disseminated MAC
 - Preferred agents: azithromycin (few drug interactions), clarithromycin

<https://clinicalinfo.hiv.gov/en/guidelines/hiv-clinical-guidelines-adult-and-adolescent-opportunistic-infections/disseminated?view=full>

37

HIV Associated Opportunistic Infections: Part 2

Immune Reconstitution Inflammatory Syndrome (IRIS)



38

Immune Reconstitution Inflammatory Syndrome (IRIS)

- **Definition:** Worsening manifestations or abrupt/atypical presentation of infection or tumor when ART started
 - Paradoxical: exacerbation of pre-existing infection or tumor
 - Unmasking: exacerbation of previously occult infection/tumor

39

Immune Reconstitution Inflammatory Syndrome (IRIS)

- **Predictors**
 - Pre therapy low CD4 cell count or high HIV RNA
 - Prior OI or recent initiation of therapy for OI
 - High pathogen load
- **Clinical Features**
 - Characterized by fevers and worsening of the underlying OI or tumor
 - May "unmask" disease at previously unrecognized site or lead to paradoxical worsening of known OI
 - Usually occurs 4-8 weeks after ART initiation; may manifest earlier or later

40

Pathogens Commonly Associated with IRIS

- Mycobacterium avium complex
- Mycobacterium tuberculosis
- Cryptococcus neoformans
- Reported with virtually all opportunistic infections and tumors

41

Mycobacterial IRIS

PATHOGEN	TYPICAL/CHARACTERISTICS OF THE DISEASE
Mycobacterium tuberculosis	<ul style="list-style-type: none"> • Worsening lung infiltrates, lymphadenitis, CNS tuberculomas
MAC	<ul style="list-style-type: none"> • Lymphadenitis; pulmonary and abdominal disease. • Bacteremia generally absent. • Elevated alkaline phosphatase may be predictive. • Severe forms of MAC IRIS with hemophagocytic lymphohistiocytosis phenotype may occur

Cecil Medicine Textbook (French and Meintjes)
<https://clinicalinfo.hiv.gov/en/guidelines/hiv-clinical-guidelines-adult-and-adolescent-opportunistic-infections/disseminated?view=full>

42

Examples of IRIS

PATHOGEN	TYPICAL/CHARACTERISTICS OF THE DISEASE
Cryptococcus neoformans	Worsening meningitis (may have brisk CSF pleocytosis)
Pneumocystis jiroveci	Exacerbation of pneumonia
Cytomegalovirus (CMV)	Vitritis
JC polyomavirus/PML	Worsening white matter changes; enhancement, edema
Human herpesvirus 8/Kaposi Sarcoma	Rapid progression of existing and/or new KS lesions
Varicella-zoster virus	Dermatomal or multidermatomal zoster; rarely myelitis

Cecil Medicine Textbook (French and Meintjes)
<https://clinicalinfo.hiv.gov/en/guidelines/hiv-clinical-guidelines-adult-and-adolescent-opportunistic-infections/disseminated?view=full>

43


Immune Reconstitution Inflammatory Syndrome (Mycobacterium avium complex)



Sereti I, IAS USA Topics in Antiviral Medicine, 2019

44

MAC IRIS in Patient with HIV



Sereti I, IAS USA Topics in Antiviral Medicine, 2019

Management of IRIS

- **Reassess Diagnosis**
 - Evaluate for concurrent, additional OIs and tumors
- **Treat IRIS**
 - Continue ART
 - Continue treatment of identified pathogen
 - NSAIDS or Corticosteroids

Summary

Multiple causes of brain lesions in people with advanced HIV; response to empiric therapy makes dx of toxoplasma encephalitis

New guidelines for induction, consolidation and maintenance therapy for cryptococcal meningitis; deferring ART for about 2-4 weeks appropriate

TB reactivation may occur even when CD4 count >200; MAC Prophylaxis no longer recommended when ART started quickly

Immune Reconstitution Inflammatory Syndrome may occur after almost all opportunistic infections or tumors: paradoxical worsening or unmasking of subclinical disease

45

46

47