



Hospital Epidemiology

Michael Klompas, MD, MPH, FIDSA, FSHEA

Professor, Harvard Medical School
Infection Control, Brigham and Women's Hospital, Boston, MA



Disclosures of Financial Relationships with Relevant Commercial Interests

- Grant Funding
 - Centers for Disease Control and Prevention
 - Agency for Healthcare Research and Quality
 - Mass Department of Public Health
- Royalties
 - UpToDate

Question #1

A surgical colleague calls you because 2 of his patients developed *Candida albicans* surgical site infections following spine surgery. You review the hospital's microbiology records and confirm that this is very unusual.

What are potential sources for this cluster?

- A. Scrub nurse wearing artificial nails
- B. Prolonged use of urinary catheters
- C. Contamination of surgical instruments
- D. Failure of peri-operative blood glucose control
- E. Use of broad-spectrum antibiotics for peri-operative prophylaxis

Nail Add-Ons & Blemishes Can Harbor Pathogens



- Nail add-ons can act as reservoirs for potentially pathogenic organisms; can persist despite cleaning with an antiseptic
- Multiple clusters linked to healthcare workers with artificial nails & infected nails
 - NICU patients with ESBL *Klebs pneumoniae* infections
 - Serratia bloodstream infections in dialysis patients linked to RN opening heparin vials with fake nails
 - NICU patients with *Pseudomonas* infections linked to healthcare workers with artificial & infected nails
 - Laminectomy surgical site infections with *Candida albicans* traced to scrub tech with artificial nails
 - Sternal wound infections with *Pseudomonas* traced to OR nurse with onychomycosis
 - Sternal wound infections with *Pseudomonas* traced to cardiac surgeon with onychomycosis

etsy.com/dk-en/listing/598625946/nurse-nail-art-decals

SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

Gupta, ICHE 2004;25:210-215
Gordon, ICHE 2007;28:743-744
Foca, N Engl J Med 2000;343:695-700

Parry, Clin Infect Dis 2001;32:352-7
McNeil, Clin Infect Dis 2001;33:317-323
Mermel, ICHE 2003;24:749-52



Essential Hand Hygiene Practices



Promote healthy hand skin & fingernails

- Fingernails should be short, healthy, and natural

- Perform hand hygiene per the WHO's **Five Moments**

- Before touching patient
- Before clean procedure
- After touching patient
- After touching body fluids
- After touching the patient's environment

- Alcohol-based hand rub typically preferred over soap & water

- Facilitate primary and secondary prevention of dermatitis


Ensure hand hygiene supplies are always readily accessible

- Widespread, convenient alcohol-based hand rub dispensers

SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

Infection Control & Hospital Epidemiology 2023;44:355-376

Better Hand Hygiene Fewer Healthcare Associated Infections




SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION


Emerg Infect Dis 2016;9:1628-1630

Cleaning Equipment Prevents Infections

Stepped-wedge cluster randomized trial of enhanced vs routine equipment cleaning, 10 units, 1 hospital, Australia
Enhanced cleaning included 3 extra hours/day of shared medical equipment cleaning + audit & feedback




SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION



Browne, Lancet Infect Dis 2024;12:1347-1356

Added Value of Gowns & Gloves Unclear

Cluster randomized cross-over trial of hand hygiene alone vs hand hygiene, gloves, and gowns when caring for infants colonized with gram negative bacilli resistant to 3rd generation cephalosporins, 12 German NICUs, 2020-2023. Cross-over @ 12months.



Faust, JAMA Network Open 2026;9(5):e2612759

Question #2

A 43-year-old man is brought to the hospital after being found unconscious with vomit in his mouth. His airway was suctioned, he was intubated for airway protection, and then transferred to the ICU. An LP was performed. Gram stain showed gram negative diplococci.

Which healthcare workers should be offered post-exposure prophylaxis?

- A. The scribe who documented the patient's emergency care
- B. The respiratory therapist that suctioned the patient's vomitus
- C. The medicine intern that did an admission physical in the ICU
- D. The radiology technician that did a portable chest x-ray in the ED
- E. The nurse that placed his IV in the ED (difficult stick, 3 attempts)

Neisseria Transmission to Healthcare Workers

Comprehensive search for occupational *Neisseria* infections in healthcare workers in England and Wales 1982-1996

Case 1	Case 2	Case 3
Provider: Doctor	Provider: EMS worker	Provider: Nurse
Full clinical exam of 9 yo with meningitis, including fundoscopy during which patient coughed into doctor's face	Transported 16 yo with meningitis to hospital. Care included airway insertion and delivery of oxygen while patient seizing in the ambulance	Nursed a 7mo with sepsis while baby being prepared for transfer to referral hospital; in close contact while child crying and coughing for at least 5h
0.5-2h contact time	0.5-2h contact time	5-6h contact time
Incubation period: 4d	Incubation period: 7d	Incubation period: 5d

Estimated 0.8 infections per 100,000 healthcare worker contacts with meningococcal patients

SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

Gillmore, Lancet 2000;356:1654-1655

Antimicrobial Prophylaxis for *Neisseria meningitidis*

Indicated for close contacts of patients with invasive disease*

- Household members (risk: 4 in 1000)
- Childcare center contacts
- Anyone directly exposed to patient's oral secretions
 - Kissing, mouth-to-mouth resuscitation
 - Endotracheal intubation, suctioning oral secretions without respiratory protection

*not indicated if *Neisseria* only isolated from sputum, nasopharynx, conjunctiva, etc.

Exposure window

- From 7 days before symptom onset through 24h after starting treatment

Prophylaxis options

- Rifampin 600mg PO q12h x 2d
- Ciprofloxacin 500mg PO x 1
- Ceftriaxone 250mg IM x 1

Cohn, MMWR Recomm Rep 2013;62(RR-2):1

Question #3


A 69-year-old man is admitted to hospital with fatigue, weight gain, and edema. He is found to have nephrotic syndrome due to multiple myeloma. On hospital day 7, a nurse notes a vesicular rash on his left flank and right chest. The patient is placed on Airborne precautions. PCR of fluid from a vesicle is positive for VZV.

Who of the following requires VariZIG?

- A. Unvaccinated seronegative nurse looking after the patient in the next room
- B. Unvaccinated seronegative respiratory therapist on rituximab for SLE
- C. Patient's pregnant nurse, 2 doses varicella vaccine as child. She is VZV IgG-
- D. Hospital roommate, 75 yo poorly controlled diabetes, unknown vax status
- E. The dermatologist that unroofed a vesicle for testing. She is VZV IgG+

Varicella Outbreak!

Cluster of 15 varicella cases, attributed to child with varicella pneumonia, Boston Children's Hospital, 1970s



SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

Leclair, NEJM 1980;302:450-453

Varicella Transmission

- **Person-to-person spread**
 - Direct contact with active lesions
 - Airborne spread from a person with respiratory involvement
 - Aerosolization from skin lesions or bedsheets (both rare but reported)
- **Incubation period:**
 - 8-21 days (usually 14-16 days)
- **Infectious period:**
 - From 24-48h before rash onset until all skin lesions crusted
- **Highly contagious if not immune:**
 - Varicella household transmission rate among susceptible individuals 85%
 - Herpes zoster household transmission rate ~25%
 - Breakthrough infections and transmissions relatively common but attenuated

Menkhaus, Lancet 1990;336:1315 (airborne spread)
Lopez, JID 2008;197:646-653 (skin lesions, linens)

Management of Varicella Exposure

- **Definition of exposure**
 - >15-60mins in same room as person with primary varicella or disseminated zoster involving the respiratory tract, or skin-to-skin contact with exposed varicella lesions
 - No exposure if HCW immune and wearing a mask or respirator
- **Management of Exposures**

Immune Status	Vaccinate?	VariZIG?	Furlough d8-21?	Monitor d8-21?
Fully vaccinated, seropositive, or prior Dx	No	No	No	Yes
Partially vaccinated	Yes	No	Depends ²	Yes
Unvaccinated & seronegative	Yes	No	Yes	Yes
Unvaccinated & unable to vaccinate ¹	No	Yes ³	Yes ⁴	Yes

¹ Vaccine contraindicated if pregnant or immunocompromised

² Furlough if vaccine was given >5d after first exposure

³ Or valacyclovir d7-13 if VariZIG not available

⁴ Furlough d8-28 if given VariZIG

SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

Question #4

A 64-year-old man with coronary disease is admitted with unstable angina and referred for urgent catheterization. A stent is placed in the circumflex. He initially improves but 3 days later develops fever, cough, and recurrent chest pain. His workup is positive for recurrent MI and influenza. The interventional cardiologist discloses that he had mild sniffles during the procedure but no fever and he wore a procedure mask at all times.


Did the cardiologist infect the patient?

- A. No, surgical masks provide excellent protection/control for respiratory viruses
- B. No, sniffles alone without fever cannot be influenza
- C. No, procedure rooms have excellent ventilation
- D. Yes, surgical masks only provide moderate protection/control for respiratory viruses
- E. Yes, surgical masks do not provide any protection against respiratory viruses

Influenza Transmission Possible with Few or No Symptoms

1,116 participants from 225 households in South Africa, tested for influenza 2x/week by PCR, 2017-2018

Source Individual Symptom Status



Influenza Transmission rate

- With 0 symptoms: 6%
- With ≥ 2 symptoms: 17%

27% of secondary flu infections were acquired from asymptomatic index cases

SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

Cohen, *Lancet Glob Health* 2021; 9:e863-874

All Respiratory Viruses are Borne by Aerosols


Viruses isolated from the air in healthcare facilities:

Influenza
RSV
SARS-CoV-2
Rhinovirus
MERS...

Wong, *Infection Control Hospital Epidemiol* 2022; PMID 35811422
Birghiana, *JAMA Network Open* 2020; 3:e2033232
Phan, *Infect Control Hospital Epidemiol* 2020; 41:259-266
Kim, *Clin Infect Dis* 2018; 66:363-9

Viral Load Predicts Transmission Risk

Secondary attack rates amongst 1,173,643 contacts of 6,263,786 index cases, UK, Jan 2021-Jan 2022



SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION


Eyre, *Lancet ID* 2023; 23:922-932

Filtration Efficiency: Surgical Mask vs. N95 Respirator

Surgical Mask with Ear Loops



Particle Penetration Behind Mask




Filtration Efficiency

38.1%

N95 Respirator



Particle Penetration Behind Mask




98.5%

Sickbert-Bennett,
JAMA Internal Med 2020;
180:1607-1612

SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

Masking and Hospital-Onset Respiratory Viral Infections


Associations between Masking Policies and Hospital-Onset Influenza, RSV, and Covid in 10 US Hospitals



SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

Pak, JAMA Network Open 2024;7(11):e2448063


Most “Aerosol Generating Procedures” Do Not Generate Aerosols



Doggett, Chest 2020; 158:2467-2473
O’Neil, Clin Infect Dis 2017;65:1342-1348
Li, Open Forum Infect Dis 2017;4(Suppl 1):S34

SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

Impact of High Flow O₂ on Respiratory Emissions



SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

Wilson, Anaesthesia 2021;76(11):1465-1474

Risk & Protection Exists on a Continuum

Factors That Increase Risk

- High community incidence
- Higher viral load
- Symptoms
- Proximity
- Longer exposure
- Poor ventilation
- Lack of masking
- Lack of vaccination

Factors That Decrease Risk

- Low community incidence
- Lower viral load
- Lack of symptoms
- Distance
- Brevity
- Good ventilation
- Mask on patient
- Mask on provider
 - N95 > KN95 > facemask
- Vaccination

Question #5


A 63-year-old man with lymphoma is admitted for chemotherapy. His course is complicated by new atrial fibrillation and hospital acquired pneumonia (treated with vancomycin, cefepime, levofloxacin). On hospital day 12 he develops severe diarrhea and is diagnosed with *C. difficile* infection.

Where did the patient most likely acquire this pathogen?

- A. From another patient on his ward (carried by healthcare workers' hands)
- B. From the previous occupant of his bed
- C. From the toilet seat of the shared bathroom in his room
- D. From the food provided by the hospital
- E. From the community (already colonized on admission)

Where Do Patients Get *C. difficile*?

Whole genome sequencing of 1,250 *C. diff* isolates from symptomatic inpatients & outpatients, Oxfordshire, UK, 2007-2011




SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

Eyre, *N Engl J Med* 2013;369:1195-1205



C. diff Colonization in ICU Patients and Progression to Infection

548 ICU patients at Johns Hopkins screened for C. difficile carriage on admission



SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

Infect Control Hospital Epidemiol 2015;36:1324-1329

So Where Do Inpatients Get C. diff From?

1. Present on admission

- Patient colonized prior to arrival, disease activates in the setting of exposure to antibiotics, antacids, immunosuppressants, and frailty

2. Transmission from symptomatic patients

- Spores carried from patient to patient via staff hands & clothing, equipment, the environment

3. Transmission from asymptomatic patients

- Spores carried from patient to patient via staff hands & clothing, equipment, the environment

Risk of C. diff Acquisition Higher if Prior Room Occupant had C. diff

Medical ICU, University of Michigan Health System, 2005-2006

Prior Room Occupant Flagged for C. diff 11.0%


Prior Room Occupant Not Flagged for C. diff 4.6%

Adjusted Hazard Ratio 2.4
(95% CI 1.2-4.5)

Infection Control Hospital Epidemiology 2011;32:201-206

Measures to Avert Hospital-Onset C. difficile Infections

Modeling study estimating the cost-effectiveness of different measures to prevent hospital-onset C. difficile infections



Barker, JAMA Network Open 2020;3(8):e2012522

Essential Practices to Prevent *C. difficile* in Hospitals



- Encourage appropriate use of antimicrobials through implementation of an antibiotic stewardship program
- Implement diagnostic stewardship to assure appropriate use and interpretation of *C. difficile* testing
 - Guide or limit use of PCR, aid in interpretation
 - Avoid testing patients if no significant diarrhea, recent positive test, or age <1 year
- Use contact precautions, single room preferred
- Adequately clean and disinfect equipment and the environment
 - Use dedicated equipment when possible (e.g., stethoscope, BP cuff, thermometer...)
- Assess the adequacy of room cleaning
 - Consider using sporicidal agents if cleaning adequate but ongoing *C. diff* transmission
- Create lab-based alerts for clinicians and infection control re new cases
- Conduct surveillance for *C. diff* infections and report to stakeholders
- Educate clinicians, enviro services, administrators, & patients about *C. difficile*
- Measure compliance with contact precautions and hand hygiene

SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION


Infection Control & Hospital Epidemiology 2023;44:527-549

Question #6

The MICU attending calls you because she's noticed 4 patients with new *Burkholderia cepacia* complex infections in her unit over the last 6 months. The patients were hospitalized during different periods. All *Burkholderia* isolates were first detected >7 days after admission.

What potential sources will you investigate?


- Are providers consistently washing their hands between patients?
- Are providers wiping down stethoscopes & phones between patients?
- Did all the patients receive care from a common healthcare worker?
- Were there any common devices amongst patients (e.g. ventilators, ECMO, bronchoscopes, ultrasound probes, etc.)?
- Did all the patients visit the same operating room?



Benjamin Franklin*



*Apocryphal. Often attributed to Franklin but unlikely he said it.

Water Avid Pathogens



- Normal inhabitants of water systems
- Promoters of persistence:
 - Biofilm forming
 - Relative resistance to disinfectants
- When clusters occur think:
 - Respiratory care equipment
 - Heating & cooling devices
 - Contaminated IV solutions & meds
 - Decorative water displays
 - Contaminated sink drains, etc.

SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION




Question #7

The CEO calls you to express her concern that ventilator-associated pneumonia rates in your hospital are double those of a competing hospital.

Which of the following measures are advised to reduce ventilator-associated pneumonia rates and improve patient outcomes?


- A. Silver coated endotracheal tubes
- B. Oral care with chlorhexidine
- C. Daily toothbrushing
- D. Placing patients in the lateral Trendelenburg position
- E. Probiotics

Circularity Between VAP Prevention Practices and the VAP Definition



SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION


Oral Care with Chlorhexidine: Significantly Lower VAP Rates



SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

Lancet Infectious Disease 2011;11:845

Oral Care with Chlorhexidine: Significantly Higher Mortality Rates




SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

BMJ 2014;348:g2197

Chlorhexidine De-Adoption Cluster Randomized Trial

Cluster randomized trial of replacing oral CHG with an oral care bundle vs continuing oral care with CHG in 6 Canadian ICUs (N=3260). Oral care bundle included twice daily toothbrushing plus mouth and lip moisturization



SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

Date, Intensive Care Med 2021;47:1295-1302

Toothbrushing: Lower Mortality, Shorter LOS

Meta-analysis of 15 randomized trials of oral care with vs without toothbrushing

	Studies	Patients	Meta-Analysis	
Hospital-acquired pneumonia*	14	2557	Risk Ratio 0.68 (95% CI 0.57-0.82)	Lower!
*12 of the 14 studies in ventilated patients				
Ventilator Days	7	1285	-1.2 days (95% CI -2.4 to -0.1)	Lower!
ICU Length of Stay	6	1284	-1.8 days (95% CI -2.9 to -0.7)	Lower!
ICU Mortality	6	1331	Risk Ratio 0.81 (95% CI 0.69-0.95)	Lower!

SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

Ehrenzeller, JAMA Internal Med 2024;184(2):131-142

Essential Practices to Prevent VAP in Adults



- Avoid intubation and prevent reintubation
 - Use high flow nasal oxygen or non-invasive positive pressure ventilation whenever safe and feasible
- Minimize sedation
 - Avoid benzodiazepines
 - Use a protocol to minimize sedation
 - Implement a ventilator liberation protocol
- Maintain and improve physical conditioning
- Elevate the head of the bed to 30-45 degrees
- Provide oral care with toothbrushing but without chlorhexidine
- Provide early enteral nutrition
- Change the ventilator circuit only if visibly soiled or malfunctioning

Infection Control & Hospital Epidemiology 2022;43:687-713

Question #8

You are part of a multidisciplinary team working to prevent central line associated bloodstream infections in your hospital. Interventions to date include education, daily patient bathing with chlorhexidine, line insertion checklists, insertion kits, and maximal sterile barrier precautions during insertion.

What additional steps should you consider implementing?

- Create a standing order for vancomycin for all patients with central lines
- Replace all central lines every 7 days
- Preferentially site all lines in the internal jugular vein whenever possible
- Require "double antiseptic" skin preparation with povidone-iodine-chlorhexidine before all insertions
- Require "double antiseptic" skin preparation with alcohol-chlorhexidine before all insertions

Essential Practices to Prevent Line Infections



Before Insertion:

- Disseminate indications for evidence-based central line use to minimize unnecessary use
- Provide education and perform competency assessments
- Daily bathing with chlorhexidine

Infection Control & Hospital Epidemiology 2022;43:553-569

Essential Practices to Prevent Line Infections



At Insertion:

- Use a checklist to assure all steps followed
- Perform hand hygiene
- Subclavian site preferred
- Use a catheter-placement kit with all necessary supplies
- Use ultrasound guidance to place the catheter
- Use maximal sterile barrier precautions
- Use an alcohol-chlorhexidine antiseptic for skin prep

Infection Control & Hospital Epidemiology 2022;43:553-569

Essential Practices to Prevent Line Infections



After Insertion:

- Ensure appropriate nurse:patient ratio and limit use of float nurses in ICUs
- Use chlorhexidine-containing dressings for central lines
- Change transparent dressings and perform site care with a chlorhexidine-based antiseptic q7d (or immediately if soiled)
- Disinfect catheter hubs, connectors, ports before each use
- Remove non-essential catheters promptly
- Replace administration sets q7d or less
- Routinely measure line infection rates and report back to unit staff & hospital leaders

Infection Control & Hospital Epidemiology 2022;43:553-569

Question #9

A 66-year-old gent with poorly controlled diabetes is admitted with fever and a swollen left knee. He underwent knee replacement 3 weeks ago. Arthrocentesis shows gram positive cocci in clusters. Culture is positive for *Staph aureus* (methicillin-susceptible). The patient is taken to the OR, the prosthesis is removed, and an antibiotic spacer is placed. The patient is devastated by the setback to his recovery and the need for more surgery.

He asks what more could have been done to prevent this infection?

- A. Obtain a urine culture before surgery to rule out occult bacteriuria
- B. Screen all patients before arthroplasty to identify *Staph aureus* carriers and decolonize them with chlorhexidine washes + nasal mupirocin
- C. Prescribe 4 weeks of antibiotic prophylaxis for all arthroplasty patients
- D. Only provide arthroplasty to patients with hemoglobin A1C's <7
- E. Ensure all knee surgeries are performed with therapeutic hypothermia

Where Do *Staph aureus* Infections Come From?

80%

Staph Bacteremia

Nasal isolates compared to blood isolates in 219 patients with *Staph aureus* bacteremia. 82% matched

von Eiff, NEJM 2001;344:11-16

of hospital acquired *Staph aureus* infections are attributable to patients' own flora (endogenous)

Surgical Site Infections


Nasal isolates compared to wound isolates in 39 patients with *Staph aureus* SSIs. 85% matched

Pert, NEJM 2002;346:1871-77

SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

Staph aureus Screening & Decolonization

917 hospitalized patients with positive *Staph aureus* nasal screens randomized to decolonization vs placebo
~90% of enrolled patients were on surgical services. Greatest benefit cardiac > ortho > vascular > GI



SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

Bode, NEJM 2010;362:9-17

Targeted vs Universal Decolonization in the ICU

REDUCE MRSA cluster-randomized trial, 74 ICUs, 43 hospitals, 74,256 patients

Screen and Isolate

Nasal MRSA screen

If positive, isolate

Screen and Decolonize

Nasal MRSA screen

If positive, isolate & decolonize with CHG baths x 5 days + mupirocin x 5 days

Universal Decolonization

No screening


Decolonize all patients with CHG baths throughout ICU stay + mupirocin x 5 days

SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

Huang et al. NEJM 2013;368:2255-65

Targeted vs. Universal Decolonization in the ICU

REDUCE MRSA cluster-randomized trial, 74 ICUs, 43 hospitals, 74,256 patients




SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

Huang et al. NEJM 2013;368:2255-65

Should We Decolonize Patients After Discharge?

1063 discharged patients colonized with MRSA randomized to education + decolonization vs education alone.
Decolonization protocol: CHG mouthwash, CHG body wash, and BID nasal mupirocin x 5d twice a month x 6mo



SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

Huang, NEJM 2019;380:638-50

Question #10


An obese 62-year-old female smoker with COPD is admitted for elective resection of adenocarcinoma of the left upper lobe. She weighs 132kg. She is intubated and undergoes left upper lobe lobectomy. Cefazolin 3g IV is administered 30mins before incision and every 4 hours during surgery. A chest tube is placed on the left side. After surgery she is admitted to the ICU for recovery.

How long should cefazolin be continued post-operatively?

- A. 0-hours – prophylaxis should be stopped after surgery
- B. 12-hours
- C. 24-hours
- D. Until the chest tube is removed
- E. Until the patient is extubated

Extending Antibiotics Beyond Surgery Does Not Prevent Infections...

Retrospective analysis of association between duration of peri-operative antibiotic prophylaxis and adverse events in 79,058 patients who underwent orthopedic, colorectal, and vascular procedures, VA hospitals, 2008-2013




SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

Branch-Elliman, JAMA Surgery 2019;154:590-598

...But Extending Antibiotics Beyond Surgery May Be Harmful

Retrospective analysis of association between duration of peri-operative antibiotic prophylaxis and adverse events in 79,058 patients who underwent orthopedic, colorectal, and vascular procedures, VA hospitals, 2008-2013



SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

Branch-Elliman, JAMA Surgery 2019;154:590-598

Essential Practices to Prevent Surgical Site Infections – Part I



- Administer antimicrobial prophylaxis according to evidence-based practices and standards
- Use parenteral and oral abx prophylaxis before colorectal surgery
- Decolonize patients with an anti-Staphylococcal agent before cardiac and orthopedic procedures (+/- those with prosthetic implants)
- Use an anti-septic vaginal prep for cesareans & hysterectomy
- Do not remove hair at the operative site (unless it interferes with surgery)
- Use skin prep containing a combination of alcohol + an antiseptic
- Maintain normothermia during perioperative period
- Use impervious plastic wound protectors for GI and biliary tract surgery
- Perform intraoperative antiseptic wound lavage
- Control blood-glucose level in the post-operative period

Infection Control & Hospital Epidemiology 2023;44:695-720

Essential Practices to Prevent Surgical Site Infections – Part II



- Perform surveillance for surgical site infections (SSIs)
- Use a checklist and/or bundle to encourage best practices
- Increase the efficiency of surveillance by utilizing automated data
- Provide ongoing SSI rate feedback to surgical and periop personnel
- Measure & provide feedback on compliance with process measures
- Educate surgeons and periop personnel about SSI prevention measures
- Educate patients and their families about SSI prevention as appropriate
- Align SSI prevention practices with evidence-based standards, rules & regulation, and manufacturers' instructions for use
- Observe and review operating room personnel and the environment of care in the operating room and central sterile reprocessing

Infection Control & Hospital Epidemiology 2023;44:695-720


Question #11

A 55-year-old woman is emergently transferred to your hospital after falling and sustaining a spinal cord injury complicated by paraplegia. She is admitted to the intensive care unit following neurosurgery.

Which of the following steps is most likely to reduce her risk of developing a catheter-associated urinary tract infection?


- Start prophylactic fosfomycin
- Screen for colonization to inform targeted antibiotic prophylaxis
- Change the urinary catheter every 7 days
- Empty the catheter drainage bag before transporting her off the unit
- Check a urinalysis daily and start pre-emptive antibiotics if she develops pyuria

Essential Practices to Prevent Catheter-Associated UTIs



SEE ENLARGED SLIDE FOLLOWING ENTIRE PRESENTATION

Infection Control & Hospital Epidemiology 2023;44:1209-1231



Question #12

A 52-year-old woman is admitted to hospital with intermittent epigastric pain. ALK, Tbili, and lipase are high. CT shows a thickened and dilated gall bladder with stones in the common bile duct. A foley is placed. The patient goes to ERCP for sphincterotomy and gallstone retrieval. Two days later she develops fever and delirium. Blood cultures are positive for carbapenem-resistant Enterobacterales.

What sources will you consider for this infection?

- A. Healthcare workers with poor hand hygiene
- B. The hospital's decorative water fountain
- C. A contaminated duodenoscope
- D. Contaminated intravenous contrast
- E. Failure to remove a foley catheter in a timely fashion

Duodenoscopes

- Notoriously difficult to sterilize
- Meta-analysis of 15 studies sampling duodenoscopes after reprocessing (925 scopes):

16% still **culture positive** after high level disinfection performed per manufacturers' instructions

9% still culture positive after **double reprocessing** or gas sterilization

- Many clusters reported



Larsen, *EClinicalMedicine* 2020;25:100451
Forbes, *JAMA Internal Med* 2023;183:191-200

Outbreak Word Associations

Pathogen	Potential Sources
Legionella	Decorative water fountains, cooling units
Pseudomonas	Respiratory care equipment, drains & sinks
Burkholderia	Water heaters & coolers (e.g., ECMO)
Carbapenem-resistant Enterobacterales	Duodenoscopes
Candida auris	Temperature probes
Mycobacterium abscessus	Ice & water machines, other water sources
Mycobacterium chimaera	Cardiac bypass heater-cooler devices
Aspergillus sp.	Construction, plants & flowers

Summary

- Equipment, hands, and clothing are commonly contaminated by bacteria
- Hand hygiene rates are inversely associated with HAI rates
- All respiratory viruses are spread by aerosols. Risk highest with high viral load, proximity, sustained exposure, poor ventilation. Surgical masks decrease risk by ~50%. N95 respirators decrease risk by ~95%
- Most aerosol generating procedures do not generate aerosols
- Most *C. difficile* is endogenous; activated during medical care in setting of antibiotics, immunosuppressants, frailty. Some hospital transmission too.
- Decolonize *Staph aureus* carriers if lines, surgery, ICU, or after discharge
- Give antibiotic prophylaxis within 60mins before incision; stop after surgery
- Contaminated water, drains, respiratory equipment, and meds can spread water-based pathogens. ICUs working on decreasing water-based care.

